

# Mathematics and numerics for data assimilation and state estimation – Lecture 4



Summer semester 2020

## Summary of lecture 3

- Probability of  $G$  given  $H$  for events  $G, H \in \mathcal{F}$ :

$$\mathbb{P}(G | H) = \frac{\mathbb{P}(G \cap H)}{\mathbb{P}(H)}$$

where we use the division-by-zero convention  $c/0 := 0$  whenever  $\mathbb{P}(H) = 0$

- Probability of  $X = a$  given  $Y$  for rv  $X, Y$ :

$$\mathbb{P}(X = a | Y)(\omega) = \mathbb{P}(X = a | \{Y = Y(\omega)\})$$

## Summary of lecture 3

- Expectation of discrete rv  $X : \Omega \rightarrow A$  given  $H \in \mathcal{F}$ :

$$\mathbb{E}[X | H] = \sum_{a \in A} a \mathbb{P}(X = a | H) = \frac{\mathbb{E}[X \mathbb{1}_H]}{\mathbb{P}(H)}$$

- Expectation of  $X$  given the rv  $Y$ :

$$\mathbb{E}[X | Y](\omega) = \mathbb{E}[X | \{Y = Y(\omega)\}]$$

- Optimal approximation property:

$$\mathbb{E}[|X - \mathbb{E}[X | Y]|^2] \leq \mathbb{E}[|X - f(Y)|^2]$$

for any mapping  $f(Y) \in \mathbb{R}^d$ .

## Last slides of lecture 3

For  $X : \Omega \rightarrow A \subset \mathbb{R}^d$  and  $Y : \Omega \rightarrow B$ , the mapping

$$g(b) := \mathbb{E}[X \mid Y = b]$$

satisfies

$$g(Y(\omega)) := \mathbb{E}[X \mid Y = Y(\omega)].$$

**Conclusion:**  $\mathbb{E}[X \mid Y]$  is an rv induced from the rv  $Y$  through the mapping  $g$ .

**Question:** Is  $\mathbb{E}[X \mid Y]$  in some sense unique?

**Question:** Given a candidate mapping  $g : B \rightarrow \mathbb{R}^d$ , is there a way to verify whether  $g(Y) = \mathbb{E}[X \mid Y]$  ?

### Definition 1 ( $\mathbb{P}$ -almost surely equal)

Two rv  $X, Y$  are said to be  $\mathbb{P}$ -almost surely equal provided

$$\mathbb{P}(\{\omega \in \Omega \mid X(\omega) = Y(\omega)\}) = 1.$$

We write

$$X = Y \quad \mathbb{P} - a.s.$$

(or just “a.s.” whenever it is clear which probability measure  $\mathbb{P}$  is considered).

Motivation:

### Example 2

$X : \Omega \rightarrow \{0, 1\}$  and  $Y : \Omega \rightarrow \{0, 1, 2\}$  with

$$\mathbb{P}(X = Y) = 1 \quad \text{and} \quad \{Y = 2\} \neq \emptyset.$$

Then  $X(\omega) \neq Y(\omega)$  for any  $\omega \in \{Y = 2\}$ , but  $X = Y$  a.s.

### Theorem 3

Consider discrete rv  $X : \Omega \rightarrow A \subset \mathbb{R}^d$  and  $Y : \Omega \rightarrow B$ . If  $g : \mathbb{R}^k \rightarrow \mathbb{R}^d$  is a mapping such that for every bounded mapping  $f : \mathbb{R}^k \rightarrow \mathbb{R}$ ,

$$\mathbb{E}[f(Y)g(Y)] = \mathbb{E}[f(Y)X] \quad (1)$$

then

$$g(Y) = \mathbb{E}[X | Y] \quad \text{a.s.}$$

**Interpretation:**  $\mathbb{E}[X | Y]$  is an a.s. unique rv of form  $g(Y)$  satisfying (1).


**Usage:** If a mapping  $B \ni b \mapsto g(b) \in \mathbb{R}^d$  satisfies (1), i.e.,

$$\sum_{b \in B} f(b)g(b)P(Y = b) = \sum_{a \in A, b \in B} f(b)aP(X = a, Y = b) \quad \forall f : B \rightarrow \mathbb{R},$$

then  $g(Y(\omega)) = \mathbb{E}[X|Y](\omega)$  for  $\mathbb{P}$ -almost all  $\omega \in \Omega$ .

## Plan for this lecture

- Properties of Random walks (steps, symmetry, recurrence)



- Convergence of random variables

## Random walks

- Are sequences of rv  $\{X_n\}$  taking values on the lattice  $\mathbb{Z}^d$  for some  $d \geq 1$ .
- The subindex  $n$  can be associated to discrete time, and  $\mathbb{Z}^d$  to discrete space (really discrete state-space).

### Definition 4 (Random walk (RW))

$X_n : \Omega \rightarrow \mathbb{Z}^d$  for  $n = 0, 1, \dots$  is an RW if the sequence of steps  $\Delta X_n := X_{n+1} - X_n$  is identically distributed and

$X_0, \Delta X_1, \Delta X_2, \dots$  are independent.




## Random walk 2

Since  $\{\Delta X_n\}$  are iid, an RW is defined by the two distributions:

- the initial state  $\mathbb{P}_{X_0}(z) = \mathbb{P}(X_0 = z)$
- the step  $\mathbb{P}_{\Delta X_0}(z) = \mathbb{P}(\Delta X_0 = z)$

### Example 5 (Simple and symmetric RW on $\mathbb{Z}^1$ )

Let  $X_0 = 0$  and  $\mathbb{P}(\Delta X_0 = \pm 1) = 1/2$ , and **let us compute**  $\mathbb{P}(X_n = k)$ .



### Solution:

Observe that the sequence  $Y_k := \mathbb{1}_{\{\Delta X_k=1\}} \sim \text{Bernoulli}(1/2)$  is iid and satisfies

$$\Delta X_k = 2Y_k - 1$$

Consequently,

$$X_n = X_0 + \sum_{k=0}^{n-1} \Delta X_k =$$

## Symmetric random walks


For rv  $X$  and  $Y$ , we introduce notation  $X \stackrel{D}{=} Y$  to say that  $X$  and  $Y$  are identically distributed.

### Definition 6 (Symmetric RW)

An RW on  $\mathbb{Z}^d$  is called symmetric if the step and the “reverse step” are identically distributed, meaning

$$X_1 - X_0 \stackrel{D}{=} X_0 - X_1.$$

**Intuition:** Equally likely to step in opposite directions.



The above RW symmetric if and only if  $p = 1/2$ .

## Simple RW

### Definition 7 (Simple RW)

An RW on  $\mathbb{Z}^d$  is called **simple** if the values of the step  $\Delta X_0$  belong to the set  $\{e_k\}_{k=1}^d$  of canonical basis vectors in  $\mathbb{R}^d$ . In other words,

$$\{X_n\} \text{ is simple} \iff \mathbb{P}(|\Delta X_0| = 1) = 1.$$

Furthermore, an RW is called **simple symmetric** if

$$\mathbb{P}(\Delta X_0 = e_k) = \mathbb{P}(\Delta X_0 = -e_k) = \frac{1}{2d}, \quad k = 1, 2, \dots, d.$$


## Example

Consider RW with steps satisfying

$$\sum_{i=1}^6 p_i = 1.$$

Constraints for the RW being

- symmetric?
- simple?
- simple symmetric?



## Matlab implementation of simple symmetric RW on $\mathbb{Z}^2$

Core idea  $X_{n+1} = X_n + \Delta X_n$  where

$$\mathbb{P}(\Delta X_n = \pm e_1) = \mathbb{P}(\Delta X_n = \pm e_2) = 1/4.$$

Use **randi(4)** in matlab to draw random integer in  $[1, 4]$ , all integers with same probability, and assign walk direction from drawn integer.

```
% Four step directions
Step = [-1 0;1 0;0 1;0 -1];

for n =1:200
    direction = randi(4);
    dX = Step(direction, :);
    plot([X(1) X(1)+dX(1)], [X(2), X(2)+dX(2)])
    X = X+dX;
end
```

See **randWalk2d.m** for more details.

## Recurrence and transience

### Definition 8

An RW on  $\mathbb{Z}^d$  with is **recurrent** if it (over its whole path  $\{X_n\}_{n \in \mathbb{N}}$ ) visits its initial state infinitely often  $\mathbb{P}$ -almost surely, and **transient** otherwise (i.e., if it visits its initial state only a finite number of times  $\mathbb{P}$ -almost surely).

- Description of a quasi-stable property: assume you are gambling, you win with probability  $\mathbb{P}(\Delta X_n = 1) = p$  lose with  $\mathbb{P}(\Delta X_n = -1) = 1 - p$ . Unless  $p = 1/2$ ,  $\{X_n\}$  is transient.
- Recurrence is a form of quasi-periodic behavior. In some settings (but not for RW) it connects spatial distribution of limit processes and time-averages over path realizations

$$\mathbb{P}(X_\infty = y) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^N \mathbb{1}_{X_n=y}.$$

## Theorem 9

Consider an RW on  $\mathbb{Z}^d$  with  $X_0 = 0$  and let

$$T := \inf\{n \geq 1 \mid X_n = 0\}$$

with the convention that  $\inf \emptyset := \infty$  and

$$N := \sum_{n \in \mathbb{N}} \mathbb{1}_{X_n=0} \quad (\text{total visits of origin})$$

Then  $\{X_n\}$  is recurrent if and only if  $\lambda := \mathbb{P}(T < \infty) = 1$  and for  $j \in \mathbb{N} \cup \{\infty\}$ ,

$$\mathbb{P}(N = j) = \begin{cases} (1 - \lambda)\lambda^{j-1} & \text{if } \lambda < 1 \\ \mathbb{1}_{j=\infty} & \text{if } \lambda = 1 \end{cases}$$

Note that  $N : \Omega \rightarrow \mathbb{N} \cup \{\infty\}$ .



## Proof of Theorem 9

Define  $\tau_0 = 0$ , and

$$\tau_{k+1} = \inf\{n > \tau_k \mid X_n = 0\} \quad \text{for } k = 0, 1, \dots$$

Note that  $\Delta\tau_k = \tau_{k+1} - \tau_k$  is a sequence of independent and  $T$ -distributed rv.

Introducing the rv

$$\bar{k} = \sup\{k \geq 0 \mid \tau_k < \infty\},$$

we can write

$$N = \sum_{n=0}^{\infty} \mathbb{1}_{X_n=0} = \sum_{k=0}^{\bar{k}} \mathbb{1}_{X_{\tau_k}=0} = \bar{k} + 1.$$

Observe that

$$\mathbb{P}(\bar{k} = j) =$$

## Which RW are recurrent?

- **(Related to FJK 2.1.13)** Symmetric and simple RW on  $\mathbb{Z}^d$  are recurrent if  $d \leq 2$  and transient otherwise.




A drunk man will eventually find his way home, but a drunk bird may get lost forever

---

*Shizuo Kakutani*

- **(Related to FJK 2.1.14)** Non-symmetric RW are always transient.



Always transient when  $p \neq 1/2$ .



# Scaling property of RW

## Theorem 10 (Random walk case of Donsker's theorem)

Let  $\{X_n\}$  be a simple symmetric RW on  $\mathbb{Z}$  with  $X_0 = 0$  and consider

$$W^{(n)}(t) := \frac{X_{\lfloor nt \rfloor}}{\sqrt{n}} \quad t \in [0, 1],$$

where  $\lfloor x \rfloor := \max\{k \in \mathbb{Z} \mid k \leq x\}$ . Then  $\{W^{(n)}(t)\}_{t \in [0,1]}$  converges in distribution to a standard Brownian motion  $\{W(t)\}_{t \in [0,1]}$ .



## Convergence of random variables

Assume you can draw iid samples  $X_k \sim \mathbb{P}_X$  and that you approximate  $\mu = \mathbb{E}[X]$  by the sample average

$$\bar{X}_M := \frac{1}{M} \sum_{k=1}^M X_k. \quad (2)$$

### Questions:

- Will  $\bar{X}_M \rightarrow \mu$  as  $M \rightarrow \infty$ , and, if so, in what sense?
- Is there a convergence rate of the form

$$\|\bar{X}_M - \mu\| \leq \frac{C}{M^\beta}$$

for some norm  $\|\cdot\|$  and some rate  $\beta > 0$ ?

## Mean-square convergence

- For rv  $Y, Z : \Omega \rightarrow \mathbb{R}^d$  we introduce the scalar product

$$\langle Y, Z \rangle_{L^2(\Omega)} := \mathbb{E}[Y \cdot Z]$$

- the function space

$$L^2(\Omega) := \{ \mathcal{F} - \text{measurable mappings } Y : \Omega \rightarrow \mathbb{R}^d \mid \mathbb{E}[|Y|^2] < \infty \}$$

with norm

$$\|Y\|_{L^2(\Omega)} := \sqrt{\mathbb{E}[|Y|^2]},$$

is a Hilbert space.

- The notation is shorthand for  $L^2(\Omega) = L^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$ .

Returning to the approximation

$$\bar{X}_M = \frac{1}{M} \sum_{k=1}^M X_k$$

- Since  $\mathbb{E}[X_k] = \mu$ , it holds that

$$\bar{X}_M - \mu = \sum_{k=1}^M \frac{X_k - \mu}{M}$$

- Since  $\{X_k - \mu\}$  is a mean-zero and independent sequence of rv, it holds for  $j \neq k$  that

$$\begin{aligned} \langle X_k - \mu, X_j - \mu \rangle_{L^2(\Omega)} &= \mathbb{E}[(X_k - \mu) \cdot (X_j - \mu)] \\ &= \sum_{(x_k, x_j) \in A \times A} (x_k - \mu) \cdot (x_j - \mu) \underbrace{\mathbb{P}(X_k = x_k, X_j = x_j)}_{=\mathbb{P}(X_k=x_k)\mathbb{P}(X_j=x_j)} \\ &= \mathbb{E}[(X_k - \mu)] \cdot \mathbb{E}[(X_j - \mu)] = 0 \end{aligned}$$

(Here we assumed discrete rv  $X_k : \Omega \rightarrow A$ , but it also holds for continuous rv.)

This yields

$$\begin{aligned}\|\bar{X}_M - \mu\|_{L^2(\Omega)}^2 &= \left\langle \sum_{k=1}^M \frac{X_k - \mu}{M}, \sum_{k=1}^M \frac{X_k - \mu}{M} \right\rangle \\ &= \end{aligned}$$

**Conclusion:** For a sequence of  $d$ -dimensional discrete independent rv  $X_i \sim \mathbb{P}_X$ ,

$$\|\bar{X}_M - \mu\|_{L^2(\Omega)} = \frac{\|X - \mu\|_{L^2(\Omega)}}{\sqrt{M}}, \quad (3)$$

i.e., the mean-square convergence rate is  $1/2$ .

## Weaker form of convergence

### Definition 11 (Convergence in probability)

A sequence of rv  $\{\bar{Y}_k\}$  converges in probability towards the rv  $Y$  if for all  $\epsilon > 0$ ,

$$\lim_{k \rightarrow \infty} \mathbb{P}(|Y_k - Y| > \epsilon) = 0.$$

### Theorem 12 (Weak law of large numbers (Durrett 2.2.14))

For a sequence of  $d$ -dimensional independent rv  $X_i \sim \mathbb{P}_X$  with  $\mathbb{E}[|X_i|] < \infty$  it holds that

$$\bar{X}_M \rightarrow \mu \quad \text{in probability.}$$



## Chebychev's inequality

To prove the theorem, we will apply **Chebychev's inequality**: for any rv  $Y$  with  $\bar{\mu} = \mathbb{E}[Y]$

$$\mathbb{P}(|Y - \bar{\mu}| > \epsilon) \leq \mathbb{E} \left[ \frac{|Y - \bar{\mu}|^2}{\epsilon^2} \right]$$


**Verification:**

**Proof of Theorem 12 (Under simplified assumption  $\mathbb{E}[|X|^2] < \infty$ )**

$$\mathbb{P}(|\bar{X}_M - \mu| > \epsilon) \leq$$

## Next time

### Discrete time and space Markov Chains



**Caption:** Quantum Cloud, designed by Antony Gormley. Random walk algorithm starting from points on the surface of an enlarged figure based on Gormley's body.