

# Mathematics and numerics for data assimilation and state estimation – Lecture 8



Summer semester 2020

## Summary of lecture 7

$$\mathbb{P}(X \in A) = \mathbb{P}_X(A), \quad A \in \mathcal{D}^d$$

$\mathcal{D}^{-1}(A)$   
 $\mathbb{R}^d$

- Random variables can be discrete, mixed or continuous.
- It is uniquely described by its distribution  $\mathbb{P}_X$ , and also by its cdf

$$F_X(x) = \mathbb{P}(X \leq x)$$

and, if it exists, also by its pdf

$$\pi_X(x) = \frac{\mathbb{P}(X \in dx)}{dx}.$$

- expectation of  $X$  given  $Y$  can be expressed through use of the
  - 1 conditional probability  $\mathbb{P}(X \in dx | Y)$  when  $Y$  is a discrete rv
  - 2 and the conditional density  $\pi_{X|Y}$  when  $X$  and  $Y$  are continuous rv.

# Overview

- 1  $L^2(\Omega)$ , sub- $\sigma$ -algebras and projections
- 2 Forward and inverse problems
  - Bayesian vs frequentist
- 3 Bayesian inversion
  - Bayesian methodology

# Overview

1  $L^2(\Omega)$ , sub- $\sigma$ -algebras and projections

2 Forward and inverse problems

- Bayesian vs frequentist

3 Bayesian inversion

- Bayesian methodology

## General definition for conditional expectation

- Mixed rv are neither discrete nor continuous.

### Example 1

$X = YZ$  where  $Y \sim \text{Bernoulli}(1/2)$  and  $Z \sim U[0, 1]$  with  $Y \perp Z$ .  
Then formally,

$$\pi_X(x) = \frac{\delta_0(x) + \mathbb{1}_{[0,1]}(x)}{2}$$

- Conditional expectations cannot always be computed using conditional densities for mixed rv.
- **Objective:** obtain a unifying definition for conditional expectations valid for all types of rv.

## Sigma algebra generated by $Y$

- For a discrete rv

$$Y(\omega) = \sum_{k=1}^{\infty} b_k \mathbb{1}_{B_k}(\omega)$$

$$B_k = Y^{-1}(b_k) \in \mathcal{F}$$

on  $(\Omega, \mathcal{F}, \mathbb{P})$ , with  $B_k = \{Y = b_k\}$  we define

$\sigma(Y) := \sigma(\{B_k\}) =$  smallest  $\sigma$ -algebra containing all events  $B_1, B_2, \dots$

- By construction  $Y$  is  $\sigma(Y)$ -measurable and  $\sigma(Y) \subset \mathcal{F}$ .
- Then, for an integrable rv  $X$ , it holds that

$$\mathbb{E}[X|Y](\omega) = \begin{cases} \frac{1}{\mathbb{P}(B_1)} \int_{B_1} X d\mathbb{P} & \text{if } \omega \in B_1 \\ \frac{1}{\mathbb{P}(B_2)} \int_{B_2} X d\mathbb{P} & \text{if } \omega \in B_2 \\ \vdots & \end{cases} = \frac{\mathbb{E}\left[\frac{\mathbb{1}_{\{B_2\}} X\right]}{\mathbb{P}(B_2)}$$

**Observations:**  $\mathbb{E}[X|Y]$  is a  $\sigma(Y)$ -measurable discrete rv for which

$$\int_B X d\mathbb{P} = \int_B \mathbb{E}[X|Y] d\mathbb{P} \quad \forall B \in \sigma(Y).$$

(hint: verify first for sets  $B_k$ , and extend to general set  $B \in \sigma(Y)$ ).

$$\omega \in B_k \Rightarrow \mathbb{E}[X|Z](\omega) = \frac{1}{P(B_k)} \int_{B_k} X d\mathbb{P} \Rightarrow \int_{B_k} (\mathbb{E}[X|Z](\omega)) d\mathbb{P} = \int_{B_k} X d\mathbb{P}$$

Seeking to preserve these properties, observe first that for

$$Y : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}^k, \mathcal{B}^k),$$

$$\sigma(Y) := \text{smallest } \sigma\text{-algebra containing } Y^{-1}(C) \quad \forall C \in \mathcal{B}^k$$

similarly satisfies  $\sigma(Y) \subset \mathcal{F}$  and that  $Y$  is  $\sigma(Y)$ -measurable.

### Definition 2 (Conditional expectation for general rv)

For rv  $X : \Omega \rightarrow \mathbb{R}^d$  and  $Y : \Omega \rightarrow \mathbb{R}^k$  defined on the same probability space, we define  $\mathbb{E}[X|Y]$  as any  $\sigma(Y)$ -measurable rv  $Z$  satisfying

$$\int_B X d\mathbb{P} = \int_B Z d\mathbb{P} \quad \forall B \in \sigma(Y).$$

## Conditioning on a $\sigma$ -algebra

One may relate  $\mathbb{E}[X | Y]$  to another kind of conditional expectation:

### Definition 3 (Expectation of $X$ given $\mathcal{V} \subset \mathcal{F}$ .)

Let  $X : \Omega \rightarrow \mathbb{R}^d$  be an integrable rv on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and assume  $\mathcal{V}$  is a  $\sigma$ -algebra  $\mathcal{V} \subset \mathcal{F}$ . Then we define  $\mathbb{E}[X | \mathcal{V}]$  as any  $\mathcal{V}$ -measurable rv  $Z$  satisfying

$$\int_B X dP = \int_B Z dP \quad \forall B \in \mathcal{V}.$$

**Observation:** Setting  $\mathcal{V} = \sigma(Y)$  implies that  $\mathbb{E}[X | Y]$  satisfies the constraints of  $\mathbb{E}[X | \sigma(Y)]$  and vice versa.

**Question:** Does  $\mathbb{E}[X | \mathcal{V}]$  exist and is it unique?

Yes,  $\mathbb{E}[X | \mathcal{V}] = \text{Proj}_{L^2(\Omega, \mathcal{V})} X$  is a.s. unique.



## Function space $L^2(\Omega, \mathcal{F})$

As an extension of  $L^2(\Omega)$  for discrete rv, we introduce the Hilbert space

$$L^2(\Omega, \mathcal{F}) = \left\{ X : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}^d, \mathcal{B}^d) \mid \int_{\Omega} |X(\omega)|^2 dP < \infty \right\}$$

$L^2(\Omega, \mathcal{F}, \mathbb{P})$   
with the scalar product

$$\langle X, Y \rangle = \int_{\Omega} X \cdot Y dP = \int_{\mathbb{R}^d \times \mathbb{R}^d} \cancel{x \cdot y} \cancel{X \cdot Y} dF(x, y)$$

and norm

$$\|X\|_{L^2(\Omega, \mathcal{F})} = \sqrt{\langle X, X \rangle}.$$

This is a Hilbert space: it is complete and for any sub- $\sigma$ -algebra  $\mathcal{V} \subset \mathcal{F}$ ,  $L^2(\Omega, \mathcal{V})$  is a closed subspace of  $L^2(\Omega, \mathcal{F})$ .

$L^2(\Omega, \mathcal{V}, \mathbb{P})$


## Orthogonal projections onto subspaces

### Definition 4

The orthogonal projection of  $X \in L^2(\Omega, \mathcal{F})$  onto the closed subspace  $L^2(\Omega, \mathcal{V})$  is defined as any rv  $Z \in L^2(\Omega, \mathcal{V})$  satisfying

$$\langle X - Z, W \rangle = 0 \quad \forall W \in L^2(\Omega, \mathcal{V}). \quad (1)$$

We write  $Z = \text{Proj}_{L^2(\Omega, \mathcal{V})} X$ .



## Orthogonal projections onto subspaces

### Definition 4

The orthogonal projection of  $X \in L^2(\Omega, \mathcal{F})$  onto the closed subspace  $L^2(\Omega, \mathcal{V})$  is defined as any rv  $Z \in L^2(\Omega, \mathcal{V})$  satisfying

$$\langle X - Z, W \rangle = 0 \quad \forall W \in L^2(\Omega, \mathcal{V}). \quad (1)$$

We write  $Z = \text{Proj}_{L^2(\Omega, \mathcal{V})} X$ .

**Exercise:** verify that  $\text{Proj}_{L^2(\Omega, \mathcal{V})} X$  satisfies the constraints of  $\mathbb{E}[X \mid \mathcal{V}]$ .

**Hint:** consider  $W = \mathbb{1}_B$  for  $B \in \mathcal{V}$

**Exercise:** verify that  $Z = \text{Proj}_{L^2(\Omega, \mathcal{V})} X$  is unique in  $L^2(\Omega, \mathcal{V})$  (and thus a.s. unique).

**Last step:** take as a fact that  $\mathbb{E}[X \mid \mathcal{V}]$  satisfies the constraint (1) of  $\text{Proj}_{L^2(\Omega, \mathcal{V})} X$ , and conclude that  $\mathbb{E}[X \mid \mathcal{V}]$  is a.s. unique.

## Summary conditional expectations

### Theorem 5

For rv  $X : \Omega \rightarrow \mathbb{R}^d$  and  $Y : \Omega \rightarrow \mathbb{R}^k$  defined on the same probability space and with  $X \in L^2(\Omega, \mathcal{F})$ , it holds that

$$\mathbb{E}[X | Y] = \mathbb{E}[X | \sigma(Y)] = \text{Proj}_{L^2(\Omega, \sigma(Y))} X.$$

# Overview

1  $L^2(\Omega)$ , sub- $\sigma$ -algebras and projections

2 Forward and inverse problems


- Bayesian vs frequentist

3 Bayesian inversion

- Bayesian methodology

# What are forward and inverse problems

## Forward problem




Possible outcomes are deduced from cause

Deductive:

General property  $\xrightarrow{\text{deduce}}$  Special case

## Inverse problem



Possible causes are induced from observations

Inductive: set of observations  $\rightarrow$  general rule

## A simple linear example

**Forward problem:** Given a matrix  $A$  and vector  $u$  compute the outcome

$$y = Au$$

**Inverse problem:** Given a matrix  $A$  and observation/effect  $y$  with either

(i)  $y \notin \text{columnSpan}(A)$ , or

(ii)  $y \in \text{columnSpan}(A)$  but  $\text{Kernel}(A) \neq \emptyset$ ,

then for (i), find the best approximate cause  $u$  to

$$Au = y$$

and for (ii), find the most suitable cause  $u$  to the above problem.

$$\|Au - y\| + \lambda \|u\|$$

# Well-posedness

## Definition 6 (J. Hadamard 1902)

A problem is called **well-posed** if

- 1 a solution exists,
- 2 the solution is unique, and
- 3 the solution is stable with respect to small perturbations in the input.

On the other hand, if any of the above conditions are not satisfied, then the problem is **ill-posed**.

**Example:** The linear forward problem

$$y = Au$$

$$y_s = Au_s$$

with fixed  $A$  and  $u$  is well-posed.

$$\left| \begin{array}{l} |y - y_s| \leq \|A\| |u - u_s| \\ \leq c |u - u_s| \end{array} \right.$$



# Well-posedness

## Definition 6 (J. Hadamard 1902)

A problem is called **well-posed** if

- 1 a solution exists,
- 2 the solution is unique, and
- 3 the solution is stable with respect to small perturbations in the input.

On the other hand, if any of the above conditions are not satisfied, then the problem is **ill-posed**.

**Example:** The inverse problem:

$$Au = y$$

with fixed  $A$  and input  $y$  is well-posed if  $A$  is invertible and  $|A^{-1}|$  is not too large. Since then for perturbed observations  $y_\delta = y + \mathcal{O}(\delta)$ ,

$$|u - u_\delta| = |A^{-1}(y - y_\delta)| \leq C|A^{-1}|\delta.$$

Otherwise, it is not a well-posed problem.

## Deterministic methods versus Bayesian inversion

We will consider extensions of the linear problem of the following form

$$Y = G(U) + \eta \quad (2)$$

where

- $Y$  is the observation
- $G$  is the possibly nonlinear forward model
- $\eta$  is a perturbation/observation noise,
- $U$  is the unknown parameter we seek to recover

**Typical deterministic approach:** View all variables as deterministic – also the perturbation. Find unique solution <sup>to</sup> an initially ill-posed problem (2) by introducing pseudo-inverse  $G^+$  and solve

$$U = G^+(Y - \eta).$$

**Bayesian approach:** View all variables as random. Model your uncertainty through input prior  $U \sim \pi_U$  and  $\eta \sim \pi_\eta$ . Solution is not a point in  $\mathbb{R}^d$ , but a posterior distribution:  $\pi_{U|Y}(\cdot|y)$  (given observation  $Y = y$ ).

## Plan for this lecture

- Bayesian methodology for solving inverse problems.
- Introduce norms to study convergence of the posterior  $\pi_{U|Y}$ .
- Well-posedness for Bayesian inversion with perturbed input model  $G_\delta \approx G$ .

## Bayesian vs frequentist

### Definition 7 (Frequentist randomness)

The probability of an event is the long-term frequency of said event occurring when repeating an experiment multiple times:

$$\mathbb{P}(X \in A) = \lim_{M \rightarrow \infty} \sum_{i=1}^M \frac{\mathbb{1}_A(X_i)}{M} \quad \text{with } X_i \sim \mathbb{P}, \quad \text{and ideally independent.}$$

- Is applicable to repeatable experiments (coin flips, card games, . . . ), and to some degree to large data experiments (elections, survey polls, etc. )
- and to some degree in settings where imaginary sampling is deduced from some form of prior information (e.g., physics argument for a coin flip being *Bernoulli*(1/2)).
- Dogmatically interpreted, not applicable to non-repeating experiments (e.g., probability that Barcelona wins a particular soccer match).

## Bayesian vs frequentist

### Definition 8 (Bayesian uncertainty)

$\mathbb{P}(X \in A \mid I)$  represents my degree of belief/confidence that  $A$  occurs given all my prior information  $I$ .

- **Constraint in definition:** If you and I have the same prior information  $I$ , then your  $\mathbb{P}(\cdot \mid I)$  should be the same as mine!
- The Bayesian approach leads to the same probability calculus as in the “usual” frequentist probability theory.
- More general than frequentist approach, as imaginary sampling really stems from a Bayesian viewpoint.
- Can assign probabilities/plausibility/belief to events which are either true or not (i.e., not at all random in the frequentist viewpoint):

$$\mathbb{P}(\text{John Doe committed the crime} \mid \text{evidence } x, y, z)$$

# Overview

- 1  $L^2(\Omega)$ , sub- $\sigma$ -algebras and projections
- 2 Forward and inverse problems
  - Bayesian vs frequentist
- 3 Bayesian inversion
  - Bayesian methodology

## Bayesian inverse problem

We consider the problem

$$Y = G(U) + \eta \quad (3)$$

where

- $U \in \mathbb{R}^d$  is the unknown parameter we seek to recover
- $Y \in \mathbb{R}^k$  is the observation
- $G : \mathbb{R}^d \rightarrow \mathbb{R}^k$  is the possibly nonlinear forward model
- $\eta$  is the observation noise

### Assumption 1

All parameters, ~~possibly~~ with the exception of  $G$  are random, described through

- $U \sim \pi_U$ ,  $\eta \sim \pi_\eta$  and  $Y \sim \pi_Y$ ,
- and  $\eta \perp U$ .

**Objective:** Given  $Y = y$  use this to improve the estimate of the first component in the joint rv  $(U, Y)$  through determining  $\pi_{U|Y}(\cdot|y)$ .

## Theorem 9 (Bayes theorem [Thm 1.2 Sanz-A., Stuart, Taeb (SST)])

Let Assumption 1 hold and assume that  $\pi_Y(y) > 0$ . Then

$$U|Y = y \sim \pi_{U|Y}(\cdot|y)$$

with

$$\pi_{U|Y}(u|y) = \frac{\pi_\eta(y - G(u))\pi_U(u)}{\pi_Y(y)}. \quad (4)$$

**Verification:** We may assume that also  $\pi_U(u) > 0$ , since otherwise (4) trivially holds. By the disintegration formula,

$$\pi_{UY}(u, y) = \pi_{U|Y}(u|y)\pi_Y(y) \quad \text{and} \quad \pi_{UY}(u, y) = \pi_{Y|U}(y|u)\pi_U(u).$$

And since  $\pi_Y(y) > 0$ , combining the above yields Bayes' rule for densities:

$$\pi_{U|Y}(u|y) = \frac{\pi_{UY}(u, y)}{\pi_Y(y)} = \frac{\pi_{Y|U}(y|u)\pi_U(u)}{\pi_Y(y)}.$$

Since  $Y|(U = u) = G(U) + \eta|(U = u) = G(u) + \eta$

$$\pi_{Y|U}(y|u) = \pi_{\eta+G(u)}(y) = \pi_\eta(y - G(u)).$$



## Remarks

likelihood function

$$\pi_{U|Y}(u|y) = \frac{\overbrace{\pi_{\eta}(y - G(u))\pi_U(u)}^{\text{likelihood function}}}{\pi_Y(y)}. \quad (5)$$

- The denominator  $\pi_Y(y)$  in (5) acts as normalizing constant is called **the evidence** or the **marginal likelihood**:

$$Z := \pi_Y(y) = \int_{\mathbb{R}^d} \pi_{\eta}(y - G(u))\pi_U(u) du.$$

- $\pi_{U|Y}(u|y)$  is the **posterior density**
- To avoid clutter, we will drop density subscripts when reference is clear ( $\pi(u) = \pi_U(u)$ ,  $\pi(u|y) = \pi_{U|Y}(u|y)$  etc.)

**Question:** Given the posterior density, how can we produce a one-value estimate of the most plausible value of  $U|Y = y$ ?

## Posterior mean and MAP estimators


### Definition 10

The posterior mean of  $U$  given  $Y = y$  is defined by

$$u_{PM} := \mathbb{E}[U|Y = y] = \int_{\mathbb{R}^d} u\pi(u|y)du$$

and the maximum a posteriori (MAP) estimator is defined by

$$u_{MAP} := \arg \max_{u \in \mathbb{R}^d} \pi(u|y).$$



## Example 11

Let  $\eta \sim N(0, \gamma^2)$  and model  $G(u) = u$  and prior

$$\pi(u) = \frac{\mathbb{1}_{(-1,1)}(u)}{2}, \quad u \sim U[-1, 1]$$


Given an observation  $Y = y$ , Bayes' theorem yields

$$\pi(u|y) = \frac{\pi_\eta(y-u)\pi_U(u)}{\pi_Y(y)} = \frac{\mathbb{1}_{(-1,1)}(u) \exp(-(y-u)^2/2\gamma^2)}{2Z}$$

with normalizing constant  $Z$ .

This yields:

$$u_{MAP} = \arg \max_{u \in \mathbb{R}} \pi(u|y) = \begin{cases} y & \text{if } y \in (-1, 1) \\ -1 & \text{if } y \leq -1 \\ 1 & \text{if } y \geq 1 \end{cases}, \quad \text{and } u_{PM} = ?$$



## Assimilating two observations

### Example 12

Consider the ordinary differential equation

$$\dot{x}(t; u) = x(t; u) \quad t > 0 \quad \text{and} \quad x(0; u) = u$$

$x(t; u) = e^t u$

and

$$G(u) = x(1; u) = ue^1,$$

and assume we have two different observations

$$Y_1 = G(U) + \eta_1, \quad \text{and} \quad Y_2 = G(U) + \eta_2,$$

with the prior density  $U \sim U[-1, 4]$ , and  $\eta_1 \sim N(0, 1)$  and  $\eta_2 \sim U[-0.5, 0.5]$  with  $\eta_1 \perp \eta_2$ .

**Problem:** Compute the posterior density for  $U | (Y_1 = 0.2, Y_2 = -0.4)$ .

## Solution to Example 12

Set  $Y = (Y_1, Y_2)$  with  $\eta = (\eta_1, \eta_2)$  and apply Theorem 1 to the joint rv  $(U, Y)$ , for the observation  $y = (0.2, -0.4)$ .

$$\begin{aligned}\pi(u|y) &= \frac{\pi_{\eta}((y_1 - G(u), y_2 - G(u)))\pi_U(u)}{Z} \\ &= \frac{\pi_{\eta_2}(y_2 - G(u))\pi_{\eta_1}((y_1 - G(u))\pi_U(u)}{Z}\end{aligned}$$

**Motivation:** For  $\pi_U(u) > 0$ ,

$$\begin{aligned}Y|(U=u) &= (G(u) + \eta_1, G(u) + \eta_2) \\ &\implies \pi(y|u) = \pi_{\eta_1\eta_2}(y_1 - G(u), y_2 - G(u)).\end{aligned}$$

and

$$\begin{aligned}\pi(u|y) &= \frac{\pi(y|u)\pi_U(u)}{Z} \\ &= \pi_{\eta_1\eta_2}(y_1 - G(u), y_2 - G(u))\pi_U(u) / Z \\ &\stackrel{\eta_1 \perp \eta_2}{=} \pi_{\eta_2}(y_2 - G(u))\pi_{\eta_1}(y_1 - G(u))\pi_U(u) / Z\end{aligned}$$

**Observation:** The posterior  $\pi(u|y_1, y_2)$  can be obtained in two ways:

- In one go:  $U|(Y_1 = y_1, Y_2 = y_2)$  mapping  $\pi_U(u)$  to  $\pi(u|y_1, y_2)$ :


$$\pi(u|y) = \frac{\pi_{\eta}((y_1 - G(u), y_2 - G(u)))\pi_U(u)}{Z}$$

- Or sequentially: 1.  $U|(Y_1 = y_1)$  mapping  $\pi_U(u)$  to  $\pi(u|y_1)$ :

$$\pi(u|y_1) = \frac{\pi_{\eta_1}(y_1 - G(u))\pi_U(u)}{Z_1}$$

and 2.  $U|(Y_1 = y_1, Y_2 = y_2)$  mapping  $\pi(u|y_1)$  to  $\pi(u|y_1, y_2)$

$$\pi(u|y_1, y_2) = \frac{\pi_{\eta_2}(y_2 - G(u))\pi(u|y_1)}{Z_2}$$



## Well-posedness for Bayesian inversion

Relation between observation and underlying parameter

$$Y = G(U) + \eta$$

$$y_j = G_{\delta} u_j + \eta$$

**Inverse problem:** what is the most likely/plausible  $U$  given  $Y = y$

**Bayesian inversion solution:** the posterior density  $\pi(u|y)$ , or a function of the density, e.g.,  $u_{PM}$  and  $u_{MAP}$ .

Hadamard's definition of well-posedness requires that a solution (i) exists, (ii) is unique and (iii) **is stable with respect to small perturbations of the input.**

By construction,  $\pi(u|y)$  exists and is unique as long as  $\pi_Y(y) > 0$ .

**Objective:** Study condition (iii) under perturbations in the model. We seek a result along the lines of

$$|G_{\delta} - G| = \mathcal{O}(\delta) \implies d(\pi^{\delta}(\cdot|y), \pi(\cdot|y)) = \mathcal{O}(\delta), \quad \text{but what is } d?$$

## Metrics on the space of pdfs

Let us introduce the space of probability density functions on  $\mathbb{R}^d$

$$\mathcal{M} := \left\{ f \in L^1(\mathbb{R}^d) \mid f \geq 0 \text{ and } \int_{\mathbb{R}^d} f(u) du = 1 \right\}$$

and recall that


$$d : \mathcal{M} \times \mathcal{M} \rightarrow [0, \infty)$$

is a metric on  $\mathcal{M}$  if for all  $\pi, \bar{\pi}, \hat{\pi} \in \mathcal{M}$

- 1  $d(\pi, \bar{\pi}) = 0 \iff \pi \stackrel{L^1}{=} \bar{\pi}$ ,
- 2  $d(\pi, \bar{\pi}) = d(\bar{\pi}, \pi)$ ,
- 3  $d(\pi, \bar{\pi}) = d(\pi, \hat{\pi}) + d(\hat{\pi}, \bar{\pi})$ .

### Definition 13 (Total variation distance)

For any  $\pi, \bar{\pi} \in \mathcal{M}$ ,


$$d_{TV}(\pi, \bar{\pi}) := \frac{1}{2} \int_{\mathbb{R}^d} |\pi(u) - \bar{\pi}(u)| du = \frac{1}{2} \|\pi - \bar{\pi}\|_{L^1(\mathbb{R}^d)}$$



## Metrics on the space of pdfs

### Definition 14 (Hellinger distance)

For any  $\pi, \bar{\pi} \in \mathcal{M}$ ,

$$d_H(\pi, \bar{\pi}) := \frac{1}{\sqrt{2}} \|\sqrt{\pi} - \sqrt{\bar{\pi}}\|_{L^2(\mathbb{R}^d)}.$$

### Lemma 15 (SST Lem 1.8)

For any  $\pi, \bar{\pi} \in \mathcal{M}$ ,

$$0 \leq d_H(\pi, \bar{\pi}) \leq 1 \quad \text{and} \quad 0 \leq d_{TV}(\pi, \bar{\pi}) \leq 1.$$

**Verification for  $d_{TV}$ :**

$$d_{TV}(\pi, \bar{\pi}) =$$

## Properties TV and Hellinger distances

### Lemma 16

For any  $\pi, \bar{\pi} \in \mathcal{M}$ ,

$$\frac{1}{\sqrt{2}} d_{TV}(\pi, \bar{\pi}) \leq d_H(\pi, \bar{\pi}) \leq \sqrt{d_{TV}(\pi, \bar{\pi})}$$

## Weak errors

The posterior mean

$$u_{PM}[\pi(\cdot|y)] = \mathbb{E}^{\pi(\cdot|y)}[u] = \int_{\mathbb{R}^d} u \pi(u|y) du$$

is one possible solution to the inverse problem.

For a perturbation in the forward model  $G_\delta = G + \mathcal{O}(\delta)$  that leads to a perturbed posterior density  $\pi^\delta(u|y)$ , we then need to bound the following to verify stability

$$|u_{PM} - u_{PM}^\delta| = |\mathbb{E}^{\pi(\cdot|y)}[u] - \mathbb{E}^{\pi^\delta(\cdot|y)}[u]|$$

More generally, for a mapping  $f : \mathbb{R}^d \rightarrow \mathbb{R}^k$ , we may be interested in bounding

$$|\mathbb{E}^{\pi(\cdot|y)}[f] - \mathbb{E}^{\pi^\delta(\cdot|y)}[f]|$$

### Lemma 17 (SST Lem 1.10)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}^k$  satisfy  $\|f\|_\infty = \sup_{u \in \mathbb{R}^d} |f(u)| < \infty$ . Then for any  $\pi, \bar{\pi} \in \mathcal{M}$ ,

$$|\mathbb{E}^\pi[f] - \mathbb{E}^{\bar{\pi}}[f]| \leq 2\|f\|_\infty d_{TV}(\pi, \bar{\pi})$$

**Verification:**

$$\begin{aligned} |\mathbb{E}^\pi[f] - \mathbb{E}^{\bar{\pi}}[f]| &= \left| \int_{\mathbb{R}^d} f(u)(\pi(u) - \bar{\pi}(u)) du \right| \\ &= \end{aligned}$$

### Lemma 18 (SST Lem 1.11)

Given  $\pi, \bar{\pi} \in \mathcal{M}$ , assume that  $f : \mathbb{R}^d \rightarrow \mathbb{R}^k$  satisfies

$$f_2^2[\pi, \bar{\pi}] := \mathbb{E}^\pi[|f(u)|^2] + \mathbb{E}^{\bar{\pi}}[|f(u)|^2] < \infty.$$

Then

$$|\mathbb{E}^\pi[f] - \mathbb{E}^{\bar{\pi}}[f]| \leq 2f_2 d_H(\pi, \bar{\pi}).$$

**Proof:**

$$|\mathbb{E}^\pi[f] - \mathbb{E}^{\bar{\pi}}[f]| = \left| \int_{\mathbb{R}^d} f(u)(\pi(u) - \bar{\pi}(u)) du \right|$$

=

Application of Lemma 18 to perturbed posterior means.

$$\begin{aligned} |u_{PM}[\pi(\cdot|y)] - u_{PM}[\pi^\delta(\cdot|y)]| &= |\mathbb{E}^{\pi(\cdot|y)}[u] - \mathbb{E}^{\pi^\delta(\cdot|y)}[u]| \\ &\leq 2f_2 d_H(\pi(\cdot|y), \pi^\delta(\cdot|y)). \end{aligned}$$

where  $f(u) = u$  for the posterior mean, and thus

$$f_2^2 = \int_{\mathbb{R}^d} |u|^2 (\pi(u|y) + \pi^\delta(u|y)) du.$$

## Next time

- Assumptions on the noise  $\eta$  and perturbations  $G_\delta$  that gives stability,
- Perturbed forward problems  $G_\delta$  to which said assumptions apply,
- Bayesian inversion in the linear setting with Gaussian distributions.