

Mathematics and numerics for data assimilation and state estimation – Lecture 9



Summer semester 2020

Overview

1 Metrics on spaces of probability density functions


2 Approximation result in $Y = G(U) + \eta$ setting

Summary of lecture 8

Conditional expectations on projections:

For rv $X : \Omega \rightarrow \mathbb{R}^d$ and $Y : \Omega \rightarrow \mathbb{R}^k$ defined on the same probability space and with $X \in L^2(\Omega, \mathcal{F})$, it holds that

$$\mathbb{E}[X | Y] = \mathbb{E}[X | \sigma(Y)] = \text{Proj}_{L^2(\Omega, \sigma(Y))} X.$$



Bayesian inversion

Inverse problem

$$Y = G(U) + \eta \quad (1)$$

- observation Y is the observation
- forward model G
- observation noise η
- U is the unknown parameter

Problem assumptions: $\eta \sim \pi_\eta$, $U \sim \pi_U$ and $\eta \perp U$.

Solution:

$$\pi_{U|Y}(u|y) = \frac{\pi_\eta(y - G(u))\pi_U(u)}{\pi_Y(y)}.$$

with $\pi_Y(y)$ often replace by equivalent normalizing constant

$$Z = Z(y) = \int \pi_\eta(y - G(u))\pi_U(u) du.$$

Definition 1 (J. Hadamard 1902)

A problem is called **well-posed** if

- 1 a solution exists,
- 2 the solution is unique, and
- 3 the solution is stable with respect to small perturbations in the input.

Objective: For the inverse problem

$$Y = G(U) + \eta,$$

study settings under which condition [3] holds for perturbations in G :

$$\underbrace{|G_\delta - G| = \mathcal{O}(\delta)}_{(i)} \implies \underbrace{d(\pi^\delta(\cdot|y), \pi(\cdot|y))}_{(ii)} = \mathcal{O}(\delta)$$

Namely, give examples where (i) holds and relate this to (ii) for different metrics.

Overview

1 Metrics on spaces of probability density functions

2 Approximation result in $Y = G(U) + \eta$ setting

Metrics on the space of pdfs

Let us introduce the space of probability density functions on \mathbb{R}^d

$$\mathcal{M} := \left\{ f \in L^1(\mathbb{R}^d) \mid f \geq 0 \text{ and } \int_{\mathbb{R}^d} f(u) du = 1 \right\}$$

and recall that

$$d : \mathcal{M} \times \mathcal{M} \rightarrow [0, \infty)$$

is a metric on \mathcal{M} if for all $\pi, \bar{\pi}, \hat{\pi} \in \mathcal{M}$

- 1 $d(\pi, \bar{\pi}) = 0 \iff \pi \stackrel{L^1}{=} \bar{\pi}$,
- 2 $d(\pi, \bar{\pi}) = d(\bar{\pi}, \pi)$,
- 3 $d(\pi, \bar{\pi}) \leq d(\pi, \hat{\pi}) + d(\hat{\pi}, \bar{\pi})$.

Definition 2 (Total variation distance)

For any $\pi, \bar{\pi} \in \mathcal{M}$,

$$d_{TV}(\pi, \bar{\pi}) := \frac{1}{2} \int_{\mathbb{R}^d} |\pi(u) - \bar{\pi}(u)| du = \frac{1}{2} \|\pi - \bar{\pi}\|_{L^1(\mathbb{R}^d)}$$

Metrics on the space of pdfs

Definition 3 (Hellinger distance)

For any $\pi, \bar{\pi} \in \mathcal{M}$,

$$d_H(\pi, \bar{\pi}) := \frac{1}{\sqrt{2}} \|\sqrt{\pi} - \sqrt{\bar{\pi}}\|_{L^2(\mathbb{R}^d)}.$$

Lemma 4 (SST Lem 1.8)

For any $\pi, \bar{\pi} \in \mathcal{M}$,

$$0 \leq d_H(\pi, \bar{\pi}) \leq 1 \quad \text{and} \quad 0 \leq d_{TV}(\pi, \bar{\pi}) \leq 1.$$

Verification for d_{TV} :

$$d_{TV}(\pi, \bar{\pi}) =$$

Properties TV and Hellinger distances

Lemma 5

For any $\pi, \bar{\pi} \in \mathcal{M}$,

$$\frac{1}{\sqrt{2}} d_{TV}(\pi, \bar{\pi}) \leq d_H(\pi, \bar{\pi}) \leq \sqrt{d_{TV}(\pi, \bar{\pi})}$$

Weak errors

The posterior mean

$$u_{PM}[\pi(\cdot|y)] = \mathbb{E}^{\pi(\cdot|y)}[u] = \int_{\mathbb{R}^d} u \pi(u|y) du$$

is one possible solution to the inverse problem.

For a perturbation in the forward model $G_\delta = G + \mathcal{O}(\delta)$ that leads to a perturbed posterior density $\pi^\delta(u|y)$, we need to bound the following to verify stability

$$|u_{PM} - u_{PM}^\delta| = |\mathbb{E}^{\pi(\cdot|y)}[u] - \mathbb{E}^{\pi^\delta(\cdot|y)}[u]|$$

More generally, for a mapping $f : \mathbb{R}^d \rightarrow \mathbb{R}^k$, we may be interested in bounding

$$|\mathbb{E}^{\pi(\cdot|y)}[f] - \mathbb{E}^{\pi^\delta(\cdot|y)}[f]|$$

Lemma 6 (SST Lem 1.10)

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}^k$ satisfy $\|f\|_{L^\infty(\mathbb{R}^d)} = \text{ess sup}_{u \in \mathbb{R}^d} |f(u)| < \infty$. Then for any $\pi, \bar{\pi} \in \mathcal{M}$,

$$|\mathbb{E}^\pi[f] - \mathbb{E}^{\bar{\pi}}[f]| \leq 2\|f\|_\infty d_{TV}(\pi, \bar{\pi})$$

Verification:

$$\begin{aligned} |\mathbb{E}^\pi[f] - \mathbb{E}^{\bar{\pi}}[f]| &= \left| \int_{\mathbb{R}^d} f(u)(\pi(u) - \bar{\pi}(u)) du \right| \\ &= \end{aligned}$$

Lemma 7 (SST Lem 1.11)

Given $\pi, \bar{\pi} \in \mathcal{M}$, assume that $f : \mathbb{R}^d \rightarrow \mathbb{R}^k$ satisfies

$$f_2^2[\pi, \bar{\pi}] := \mathbb{E}^\pi[|f(u)|^2] + \mathbb{E}^{\bar{\pi}}[|f(u)|^2] < \infty.$$

Then

$$|\mathbb{E}^\pi[f] - \mathbb{E}^{\bar{\pi}}[f]| \leq 2f_2 d_H(\pi, \bar{\pi}).$$

Proof:

$$\begin{aligned} |\mathbb{E}^\pi[f] - \mathbb{E}^{\bar{\pi}}[f]| &= \left| \int_{\mathbb{R}^d} f(u)(\pi(u) - \bar{\pi}(u)) du \right| \\ &= \end{aligned}$$

Application of Lemma 7 to perturbed posterior means.

$$\begin{aligned} |u_{PM}[\pi(\cdot|y)] - u_{PM}[\pi^\delta(\cdot|y)]| &= |\mathbb{E}^{\pi(\cdot|y)}[u] - \mathbb{E}^{\pi^\delta(\cdot|y)}[u]| \\ &\leq 2f_2 d_H(\pi(\cdot|y), \pi^\delta(\cdot|y)). \end{aligned}$$

where $f(u) = u$ for the posterior mean, and thus

$$f_2^2 = \int_{\mathbb{R}^d} |u|^2 (\pi(u|y) + \pi^\delta(u|y)) du.$$

Example 8 (Extension of MAP estimator example, Lecture 8)

Consider the problem (1) with $\eta \sim N(0, \gamma^2)$, $U \sim U[-1, 1]$, $G(u) = u$ and $G_\delta(u) = u + \delta$ for some fixed $\gamma > 0$ and $\delta > 0$.

Solutions:

$$\pi(u|y) = \frac{e^{-(y-u)^2/2\gamma^2} \mathbb{1}_{(-1,1)}(u)}{2Z(y)}$$

and


$$\pi^\delta(u|y) = \frac{e^{-(y-(u+\delta))^2/2\gamma^2} \mathbb{1}_{(-1,1)}(u)}{2Z(y-\delta)} = \pi(u|y-\delta)$$

Recalling that

$$u_{MAP}[\pi(\cdot|y)] = \arg \max_{u \in \mathbb{R}} \pi(u|y) = \begin{cases} y & \text{if } y \in (-1, 1) \\ -1 & \text{if } y \leq -1 \\ 1 & \text{if } y \geq 1 \end{cases}$$

implies that $|u_{MAP}[\pi(\cdot|y)] - u_{MAP}[\pi^\delta(\cdot|y)]| \leq \delta$.

Distance between u_{MAP} and u_{MAP}^δ when $\gamma = 1$, $y = 0.1$ and $\delta = 0.2$.



Exercise

Prove that also

$$|u_{PM}[\pi(\cdot|y)] - u_{PM}[\pi^\delta(\cdot|y)]| = \mathcal{O}(\delta).$$

Overview

1 Metrics on spaces of probability density functions

2 Approximation result in $Y = G(U) + \eta$ setting

Approximation assumptions

By introducing the notation

$$g(u) := \pi_\eta(y - G(u)) \quad \text{and} \quad g_\delta(u) := \pi_\eta(y - G_\delta(u)),$$

we have

$$\pi(u|y) = \frac{g(u)\pi_U(u)}{Z} \quad \text{and} \quad \pi^\delta(u|y) = \frac{g_\delta(u)\pi_U(u)}{Z^\delta}.$$

Assumption 1

Assume there exists constant $K_1, K_2 > 0$ such that for sufficiently small $\delta > 0$,

- (i) $\sqrt{\mathbb{E}^{\pi_U}[|\sqrt{g} - \sqrt{g_\delta}|^2]} \leq K_1\delta$
- (ii) $\|\sqrt{g}\|_{L^\infty(\mathbb{R}^d)} + \|\sqrt{g_\delta}\|_{L^\infty(\mathbb{R}^d)} \leq K_2$

Approximation results

Theorem 9

If Assumption 1 holds, then there exists $c_1, c_2, c_3 > 0$ such that for sufficiently small $\delta > 0$

$$|Z - Z^\delta| \leq c_1 \delta \quad \text{and} \quad Z, Z^\delta > c_2 \quad [\text{SST Lemma 1.15}]$$

and

$$d_H(\pi(\cdot|y), \pi^\delta(\cdot|y)) \leq c_3 \delta \quad [\text{SST Theorem 1.14}]$$

where we recall that

$$d_H(\pi, \bar{\pi}) = \frac{1}{\sqrt{2}} \|\sqrt{\pi} - \sqrt{\bar{\pi}}\|_{L^2}.$$

Proof idea Lemma 1.15

$$|Z - Z^\delta| = \left| \int (g(u) - g_\delta(u)) \pi_U(u) du \right|$$

Positivity: $Z = \pi_Y(y) > 0$ by assumption, so by ...

Proof idea Thm 1.14

$$\begin{aligned}d_H(\pi(\cdot|y), \pi^\delta(\cdot|y)) &= \frac{1}{\sqrt{2}} \|\sqrt{\pi} - \sqrt{\pi^\delta}\|_2 \\&= \frac{1}{\sqrt{2}} \left\| \sqrt{\frac{g\pi_U}{Z}} - \sqrt{\frac{g_\delta\pi_U}{Z^\delta}} \right\|_2 \\&\leq \frac{1}{\sqrt{2}} \left\| \sqrt{\frac{g\pi_U}{Z}} - \sqrt{\frac{g_\delta\pi_U}{Z}} \right\|_2 + \frac{1}{\sqrt{2}} \left\| \sqrt{\frac{g_\delta\pi_U}{Z}} - \sqrt{\frac{g_\delta\pi_U}{Z^\delta}} \right\|_2 \\&\leq\end{aligned}$$

Summary of well-posedness result

Recall that

$$g(u) := \pi_Y(y - G(u)) \quad \text{and} \quad g_\delta(u) := \pi_Y(y - G_\delta(u)),$$

which yields

$$\pi(u|y) = \frac{g(u)\pi_U(u)}{Z} \quad \text{and} \quad \pi^\delta(u|y) = \frac{g_\delta(u)\pi_U(u)}{Z^\delta}.$$

Summary results: If for sufficiently small $\delta > 0$

- (i) $\|\sqrt{g} - \sqrt{g_\delta}\|_{L^2(\mathbb{R}^d)} = \mathcal{O}(\delta)$
- (ii) $\|\sqrt{g}\|_{L^\infty(\mathbb{R}^d)} + \|\sqrt{g_\delta}\|_{L^\infty(\mathbb{R}^d)} < \infty$

Then the well-posedness condition [3] holds in the following sense:

$$d_H(\pi(\cdot|y), \pi^\delta(\cdot|y)) = \mathcal{O}(\delta).$$

Example with unspecified model where (i) and (ii) hold

Consider setting where $\|G_\delta - G\|_\infty = \mathcal{O}(\delta)$

$$\|G\|_\infty + \|G_\delta\|_\infty < \infty \quad \text{and} \quad \eta \sim N(0, 1).$$

Then

$$\begin{aligned} \sqrt{g(u)} - \sqrt{g_\delta(u)} &= \sqrt{\pi_\eta(y - G(u))} - \sqrt{\pi_\eta(y - G_\delta(u))} \\ &= \frac{1}{(2\pi)^{1/4}} \left(\exp\left(\frac{-(y - G(u))^2}{4}\right) - \exp\left(\frac{-(y - G_\delta(u))^2}{4}\right) \right) \end{aligned}$$

\leq

$$= \mathcal{O}(\delta).$$

$$\text{and} \quad \|\sqrt{g}\|_\infty = \|\sqrt{g_\delta}\|_\infty = \frac{1}{(2\pi)^{1/4}}.$$

And a specified model which may lead to stability

Consider the ordinary differential equation

$$\dot{x}(t; u) = x(t; u) \quad t > 0 \quad \text{and} \quad x(0; u) = u \quad \text{for } u \in [-1, 1],$$

and the associated explicit-Euler numerical solution

$$X_{n+1}^\delta = X_n^\delta(1 + \delta), \quad X_0^\delta = u.$$

The forward model is the solution flow map from $t = 0$ to $t = 1$:

$$G(u) = x(1; u) = ue^1 \quad \text{and} \quad G_\delta(u) = X_{\lfloor \delta^{-1} \rfloor}^\delta(1 + (1 - \delta \lfloor \delta^{-1} \rfloor)).$$

For simplicity, we assume that $\delta^{-1} = N \in \mathbb{N}$. Then $G_\delta(u) = X_N^\delta$.

For $t_k = k\delta$, and note that

$$X(t_{k+1}) = e^\delta X(t_k).$$

For $E_k := |X(t_k) - X_k^\delta|$ it then holds that

$$E_{k+1} \leq (e^\delta - (1 + \delta))|X(t_k)| + (1 + \delta)E_k$$

Verification:

Consequently,

$$E_N = |G(u) - G_\delta(u)| \leq \underbrace{(e^\delta - (1 + \delta))}_{\leq c\delta^2} |X(t_{N-1})| + (1 + \delta)E_{N-1}$$

\leq

$$\leq c\delta^2 \sum_{k=0}^{N-1} (1 + \delta)^{N-1-k} |X(t_k)| + (1 + \delta)^N E_0 \leq c\delta e^1 |u| \leq c\delta.$$

For the **relevant** $u \in [-1, 1]$, we have shown that

$$\|G - G_\delta\|_{L^\infty([-1,1])} \leq c\delta,$$

where $c > 0$ satisfies

$$|e^\delta - (1 + \delta)| \leq c\delta^2 \quad \forall \delta \in (0, \delta^+) \quad (2)$$

Note also that

$$\|G\|_{L^\infty([-1,1])} + \|G_\delta\|_{L^\infty([-1,1])} \leq e^1 + (1 + \delta)^{1/\delta} \leq 2e^1.$$

Exercise: For any $\delta \in (0, \delta^+ = 1)$, show that $c = e^1/2$ satisfies (2).

Comments:

- Relevant u values not being the whole of \mathbb{R}^d may be motivated for instance by π_U having compact support.
- See also [SST 1.1.3] for a more general example of forward models stable under perturbations.

Next time

- Bayesian inversion in the linear-Gaussian setting,
- For the linear-Gaussian setting, study the posterior density in the small noise limit $\eta \sim N(0, \Gamma)$ when $|\Gamma| \rightarrow 0$.
- How informative is the MAP estimator?